

Design Considerations for Assessing Ground-Water Quality in a Regional Aquifer System:

EXPLANATION

MAJOR AQUIFER STUDIES

Platte River Alluvium

LAND-USE STUDIES

Irrigated Ag. - corn

Irrigated Ag. - cotton

REGIONAL TRANSECTS = =

Irrigated Ag. - corn

SPECIAL STUDIES

Public-Supply Wells

■ TANC Study, York, NE

Source-Water Assessment-GW

Arikaree Group and Brule Fm

NHP Ogallala

Quaternary

bbruce@usgs.gov kdennehy@usgs.gov

pmcmahon@usgs.gov

Bret W. Bruce¹ Kevin F. Dennehy² Peter B. McMahon¹ Jason J. Gurdak¹ Sharon L. Qi³ jjgurdak@usgs.gov slqi@usgs.gov

¹U.S. Geological Survey, Lakewood, CO ²U.S. Geological Survey, Reston, VA ³ U.S. Geological Survey, Vancouver, WA


ENVIRONMENTAL SETTING

The High Plains aquifer is an important regional system that covers about 174,000 mi² in parts of eight western states and supports one of the largest agriculture economies in the U.S. The High Plains Regional Ground Water (HPGW) study, part of the USGS National Water-Quality Assessment (NAWQA) Program, was tasked with assessing the current condition of ground-water quality in the High Plains aquifer and investigating the natural and human processes controling this condition.

The High Plains Aquifer System is comprised of six hydraulically connected hydrogeologic units of late Tertiary or Quaternary age each having a different set of ground-water characteristics. The Ogallala Formation of Miocene age is the principal geologic unit comprising about 70% of the aquifer system.

The large geographic area of the High Plains coupled with the variation across the area in the underlying bedrock, climate conditions, land- and water-use, depth to ground water, saturated thickness, and water-table declines present unique challenges to a comprehensive investigation of ground-water quality.

This poster provides visual aids for discussion of the design and implimentation of the HPGW study. This study approach might be useful as a template for other evaluations of regional water-quality




Location of the High Plains aquifer and the hydrogeologic units that comprise the aquifer system.

Average Annual Precipitation






A National Resource 27% of the nation's agriculture 30% of the nation's ground-water irrigation 14.5 million cattle 10.4 million swine 2.3 million people



High Plains Variability Regional Climate Differences Land-use distribution and rotations Depth to water and water-table changes Large range of saturated thickness Ground-water age (<10 yrs to >10,000 yrs)



Distribution of Agriculture

Depth to Ground Water

STUDY DESIGN CONSIDERATIONS: A Holistic Approach

Regional assessment of water quality in the High Plains aquifer required taking into account the large geographic extent of the aquifer and the spatial variation in lithology, depth to water, saturated thickness, and ground-water age. Regional differences in climate, water-use, land-use, and crop types, also influenced the location of process related studies and the choice of analytical methods used for water-quality analysis. Areally and vertically nested measurements of ground-water quality coupled with focused studies of processes controlling water movement, chemical transport, and chemical transformation were used to provide comparative results that could later be "scaled up" to a regional understanding of the entire system.

Primary Local Concerns (from survey) Nutrient contamination of ground water from CAFOs Effects of saline ground water from bedrock Effects of urban and agricultural land-use on water quality Impacts to shallow alluvial aquifers from use of surface water Effects of focused recharge

Design Considerations Large area/Limited funding Multiple hydrogeologic units Large spatial, vertical, and temporal variation Multiple scales of assessment Phased logistics and integration of results


The High Plains Regional Ground-Water (HPGW) study was designed around a series of broad-scale and process oriented studies, consistently applied in a regionally phased approach, that proved to be logistically and financially manageable; evaluated most of the primary hydrogeologic, landuse, and climatic settings; addressed local differences; and permitted scaling of results to meet interpretive needs. Primary study components were:


Major Aquifer Studies: existing domestic wells, broad suite of chemical constituents, assesses regional water quality conditions Land-Use Studies: installed water-table monitoring wells, analytes indicative of associated land use, determines impacts of dominant land-use settings on ground-water quality

Regional Transect Studies: vertically nested monitoring wells, selective analytes and range of age dating techniques, identifies vertical gradients in ground-water chemistry and age (includes effects of underlying bedrock)

Unsaturated-Zone Studies: various instrumentation between land surface and water table, continuous monitoring of moisture and periodic monitoring of chemistry, understanding of water movement (recharge) and chemical loading to aquifer

Phased Approach by Sub-Region; Areal Nesting of Studies






Phased approach lowered annual cost Assessed both regional gradients and local variability Resulted in comparable data between sub-regions Nested process-related studies provide understanding of controls on regional ground-water quality


Regionally Comparable Data; Vertical Nesting of Studies





Even distribution of studies Investigates water/chemical flux from land surface to water table Assesses vertical differences in ground-water quality Integrates multiple time scales Evaluates affect of underlying bedrock

IMPLEMENTATION OF WATER-QUALITY ASSESSMENT





Unsaturated-zone study installation

More than 300 domestic wells About 120 monitoring wells Nine unsaturated-zone study sites Three regional transect studies

Benefits of Design

Regionally Comparable Data Sets

Addresses Variability in Controlling Factors


Regional Differences Observed

Provides Understanding at Multiple Scales

Holistic Assessment of Chemical Lifecycle (Source-Transport-Receptor Model)

Revised Conceptual Model





Interpretation using S-T-R model

Future Directions

Integrate with remote sensing


Determine climatic controls on aquifer recharge

Refine techniques for data extrapolation

Update ground-water flow model

Couple climate-change, ground-water flow, and agricultural economic models

For Updates, Data, Publications, and Future Results: http://co.water.usgs.gov/nawqa/hpgw/HPGW_home.html



Weather station located at rangeland unsaturated zone monitoring site.